



University of Kentucky  
UKnowledge

Plant Pathology Faculty Publications

Plant Pathology

1-1-2018

# ICTV Virus Taxonomy Profile: *Partitiviridae*

Eeva J. Vainio

*Natural Resources Institute Finland, Finland*

Sotaro Chiba

*Nagoya University, Japan*

Said A. Ghabrial

*University of Kentucky, [saghab00@uky.edu](mailto:saghab00@uky.edu)*

Edgar Maiss

*Leibniz University Hannover, Germany*

Marilyn Roossinck

*Pennsylvania State University*

*See next page for additional authors*

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**

Follow this and additional works at: [https://uknowledge.uky.edu/plantpath\\_facpub](https://uknowledge.uky.edu/plantpath_facpub)



Part of the [Genetics and Genomics Commons](#), [Virology Commons](#), and the [Viruses Commons](#)

## Repository Citation

Vainio, Eeva J.; Chiba, Sotaro; Ghabrial, Said A.; Maiss, Edgar; Roossinck, Marilyn; Sabanadzovic, Sead; Suzuki, Nobuhiro; Xie, Jiatao; Nibert, Max; and ICTV Report Consortium, "ICTV Virus Taxonomy Profile: *Partitiviridae*" (2018). *Plant Pathology Faculty Publications*. 83.

[https://uknowledge.uky.edu/plantpath\\_facpub/83](https://uknowledge.uky.edu/plantpath_facpub/83)

This Article is brought to you for free and open access by the Plant Pathology at UKnowledge. It has been accepted for inclusion in Plant Pathology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**Authors**

Eeva J. Vainio, Sotaro Chiba, Said A. Ghabrial, Edgar Maiss, Marilyn Roossinck, Sead Sabanadzovic, Nobuhiro Suzuki, Jiatao Xie, Max Nibert, and ICTV Report Consortium

**ICTV Virus Taxonomy Profile:** *Partitiviridae*

**Notes/Citation Information**

Published in *Journal of General Virology*, v. 99, issue 1, p. 17-18.

This is an open access article published by the Microbiology Society under the [Creative Commons Attribution License](#).

**Digital Object Identifier (DOI)**

<https://doi.org/10.1099/jgv.0.000985>



# ICTV Virus Taxonomy Profile: *Partitiviridae*

Eeva J. Vainio,<sup>1,\*</sup> Sotaro Chiba,<sup>2</sup> Said A. Ghabrial,<sup>3</sup> Edgar Maiss,<sup>4</sup> Marilyn Roossinck,<sup>5</sup> Sead Sabanadzovic,<sup>6</sup> Nobuhiro Suzuki,<sup>7</sup> Jiatao Xie,<sup>8</sup> Max Nibert<sup>9</sup> and ICTV Report Consortium

## Abstract

The *Partitiviridae* is a family of small, isometric, non-enveloped viruses with bisegmented double-stranded (ds) RNA genomes of 3–4.8 kbp. The two genome segments are individually encapsidated. The family has five genera, with characteristic hosts for members of each genus: either plants or fungi for genera *Alphapartitivirus* and *Betapartitivirus*, fungi for genus *Gammapartitivirus*, plants for genus *Deltapartitivirus* and protozoa for genus *Cryspovirus*. Partitiviruses are transmitted intracellularly via seeds (plants), oocysts (protozoa) or hyphal anastomosis, cell division and sporogenesis (fungi); there are no known natural vectors. This is a summary of the International Committee on Taxonomy of Viruses (ICTV) Report on the taxonomy of the *Partitiviridae*, which is available at [www.ictv.global/report/partitiviridae](http://www.ictv.global/report/partitiviridae).

Table 1. Characteristics of the family *Partitiviridae*

Typical member:	<i>Atkinsonella hypoxylon virus</i> , 2H (RNA1, L39125; RNA2, L39126), species <i>Atkinsonella hypoxylon virus</i> , genus <i>Betapartitivirus</i>
Genome	3–4.8 kbp of linear bisegmented dsRNA
Virion	Isometric, non-enveloped, 25–43 nm in diameter; dsRNA1 and dsRNA2 are separately encapsidated
Replication	Cytoplasmic. Genomic RNA acts as a template for mRNA synthesis within the virus particle; transcription occurs by a semiconservative mechanism
Translation	From monocistronic positive-sense transcripts of both genomic dsRNAs
Host range	Plants, fungi and protozoa
Taxonomy	Five genera, including >40 species, and 15 species unassigned to a genus

## VIRION

Virus particles are isometric, non-enveloped, and 25–43 nm in diameter (Table 1, Fig. 1a, b). Each capsid is composed of 120 copies of a single protein arranged as 60 dimers with *T*=1 icosahedral symmetry [1]. Dimeric surface protrusions are frequently observed on viral capsids. One or two molecules of RNA-dependent RNA polymerase (RdRP) are packaged inside each particle [2].

## REPLICATION

Each dsRNA is monocistronic. The RdRP is believed to function as both a transcriptase and a replicase and catalyzes *in vitro* end-to-end transcription of each dsRNA to produce mRNA by a semi-conservative mechanism. Virions accumulate in the cytoplasm.

## GENOME

Members of all five genera possess two essential genome segments, dsRNA1 and dsRNA2, each containing one large ORF on the positive-strand RNA molecule (Fig. 2). The smaller of the two dsRNA genome segments usually encodes the coat protein (CP) and the larger usually encodes the virion-associated RNA polymerase. The linear dsRNA segments are separately encapsidated. Additional dsRNA segments (satellite or defective) may also be present.

## TAXONOMY

### *Alphapartitivirus*

Members of the genus *Alphapartitivirus* infect either plants, or ascomycetous or basidiomycetous fungi. The two


Received 21 November 2017; Accepted 23 November 2017

**Author affiliations:** <sup>1</sup>Natural Resources Institute Finland (Luke), Helsinki 00790, Finland; <sup>2</sup>Asian Satellite Campuses Institute, Nagoya University, Nagoya 464-0861, Japan; <sup>3</sup>Department of Plant Pathology, University of Kentucky, Lexington, KY 40546, USA; <sup>4</sup>Institute of Horticultural Production Systems, Leibniz University Hannover, Hannover 30419, Germany; <sup>5</sup>Center for Infectious Disease Dynamics, Pennsylvania State University, University Park, PA 16802, USA; <sup>6</sup>Department of Biochemistry, Molecular Biology, Entomology and Plant Pathology, Mississippi State University, MS 39762, USA; <sup>7</sup>Institute of Plant Science and Resources, Okayama University, Chuo 2-20-1, Kurashiki, 710-0046, Japan; <sup>8</sup>College of Plant Science and Technology, Huazhong Agricultural University, Wuhan 430070, Hubei Province, PR China; <sup>9</sup>Department of Microbiology and Immunobiology, Harvard Medical School, Boston, MA 02115, USA.

**\*Correspondence:** Eeva J. Vainio, [eeva.vainio@luke.fi](mailto:eeva.vainio@luke.fi)

**Keywords:** *Partitiviridae*; ICTV; taxonomy; *Alphapartitivirus*; *Betapartitivirus*; *Deltapartitivirus*; *Gammapartitivirus*; *Cryspovirus*.

**Abbreviations:** CP, coat protein; RdRP, RNA-dependent RNA polymerase.



**Fig. 1.** (a) Transmission electron micrograph of negatively-stained purified particles of *Penicillium stoloniferum* virus S, a representative member of the genus *Gammapartitivirus*. (b) Cryo-EM reconstructions of *Penicillium stoloniferum* virus S at 0.45 nm resolution, and rendered with radial colour mapping.


essential dsRNA genome segments are individually about 1.9–2.0 kbp (dsRNA1) and 1.7–1.9 kbp (dsRNA2), typically containing a poly(A) tract near the plus-strand 3′-terminus. There is a single major CP with predicted *Mr* of 51–57 kDa. Plant alphapartitiviruses cause persistent infections, whereas some fungal alphapartitiviruses cause host effects, such as hypovirulence or a reduced growth rate [3, 4].

### **Betapartitivirus**

Members of the genus *Betapartitivirus* infect either plants, or ascomycetous or basidiomycetous fungi. The two essential dsRNA genome segments are about 2.2–2.4 kbp (dsRNA1) and 2.1–2.4 kbp (dsRNA2), typically containing a poly(A) tract near the plus-strand 3′-terminus. There is a single major CP with predicted *Mr* of 71–77 kDa. Plant betapartitiviruses cause persistent infections [5, 6]. Some fungal betapartitiviruses cause reduced host virulence and changes in colony morphology [7].

### **Gammapartitivirus**

All known members of the genus *Gammapartitivirus* infect ascomycetous fungi. The two essential dsRNA segments are about 1.6–1.8 kbp (dsRNA1) and 1.4–1.6 kbp (dsRNA2). There is a single major CP with predicted *Mr* of 44–47 kDa. Most gammapartitiviruses seem to induce latent infections. *Aspergillus fumigatus* partitivirus 1, a related, unclassified virus, has been associated with host effects.



**Fig. 2.** *Atkinsonella hypoxylon* virus [10], an isolate of the type species of the genus *Betapartitivirus*, has a bipartite genome consisting of dsRNA1 and dsRNA2.

### **Deltapartitivirus**

All known members of the genus *Deltapartitivirus* induce persistent infections in plants [8]. They are transmitted by ovule and pollen to the seed embryo. The two essential dsRNA segments are individually 1.6–1.7 kbp (dsRNA1) and 1.4–1.6 kbp (dsRNA2). There is a single major CP with predicted *Mr* of 38–49 kDa.

### **Cryspovirus**

Members of the genus *Cryspovirus* infect apicomplexan protozoa of the genus *Cryptosporidium* [9]. The viral genome comprises two dsRNA segments, which are individually 1.5 and 1.8 kbp. There is a single major CP with predicted *Mr* of 37 kDa. Virions are disseminated within *Cryptosporidium* oocysts. Infections of the *Cryptosporidium* host cells appear to be latent.

## **RESOURCES**

Full ICTV Online (10th) Report:

[www.ictv.global/report/partitiviridae](http://www.ictv.global/report/partitiviridae).

### **Funding information**

Production of this summary, the online chapter, and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA).

### **Acknowledgements**

Members of the ICTV Report Consortium are Elliot J. Lefkowitz, Andrew J. Davison, Stuart G. Siddell, Sead Sabanadzovic, Donald B. Smith, Richard J. Orton and Peter Simmonds.

### **Conflicts of interest**

The authors declare that there are no conflicts of interest.

### **References**

1. Nibert ML, Tang J, Xie J, Collier AM, Ghabrial SA *et al*. 3D structures of fungal partitiviruses. *Adv Virus Res* 2013;86:59–85.
2. Nibert ML, Ghabrial SA, Maiss E, Lesker T, Vainio EJ *et al*. Taxonomic reorganization of family *Partitiviridae* and other recent progress in partitivirus research. *Virus Res* 2014;188: 128–141.
3. Chiba S, Lin YH, Kondo H, Kanematsu S, Suzuki N. Effects of defective interfering RNA on symptom induction by, and replication of, a novel partitivirus from a phytopathogenic fungus, *Rosellinia necatrix*. *J Virol* 2013;87:2330–2341.
4. Vainio EJ, Hantula J. Taxonomy, biogeography and importance of *Heterobasidium* viruses. *Virus Res* 2016;219:2–10.
5. Lesker T, Rabenstein F, Maiss E. Molecular characterization of five betacryptoviruses infecting four clover species and dill. *Arch Virol* 2013;158:1943–1952.
6. Roossinck MJ. Lifestyles of plant viruses. *Philos Trans R Soc Lond B Biol Sci* 2010;365:1899–1905.
7. Xiao X, Cheng J, Tang J, Fu Y, Jiang D *et al*. A novel partitivirus that confers hypovirulence on plant pathogenic fungi. *J Virol* 2014; 88:10120–10133.
8. Sabanadzovic S, Valverde RA. Properties and detection of two cryptoviruses from pepper (*Capsicum annuum*). *Virus Genes* 2011; 43:307–312.
9. Nibert ML, Woods KM, Upton SJ, Ghabrial SA. *Cryspovirus*: a new genus of protozoan viruses in the family *Partitiviridae*. *Arch Virol* 2009;154:1959–1965.
10. Oh CS, Hillman BI. Genome organization of a partitivirus from the filamentous ascomycete *Atkinsonella hypoxylon*. *J Gen Virol* 1995; 76:1461–1470.